

The dosage of **ALTOSID® Premix** is proportional to body weight. The recommended consumption of this product is 0.2 oz. (0.0125 lb.) to 0.4 oz. (0.025 lb.) per 100 lbs. of body weight per month (30 days). Feed by either of the following methods.

MIXED WITH FREE-CHOICE-FED MINERALS

Mix **ALTOSID Premix** with free-choice-fed minerals to produce an **ALTOSID Premix** and mineral mixture containing 0.01% to 0.02% active ingredient (S-methoprene) as shown in the following table.

0.01 % OR 4 OZ. PER DAY FEEDING RATE

ALTOSID Premix	5 lbs.	(2.44%)
Free-Choice Minerals	200 lbs.	(97.56%)
Total	205 lbs.	(100.00%)

0.02 % OR 2 OZ. PER DAY FEEDING RATE

ALTOSID Premix	5 lbs.	(4.76%)
Free-Choice Minerals	100 lbs.	(95.24%)
Total	105 lbs.	(100.00%)

Place the mixture of **ALTOSID Premix** and minerals near watering or loafing areas and provide 1 feeder for each 15 to 20 head. Put out only a 5 to 7 day supply of the mixture at one time and protect it from the rain.

When the recommended mixture of **ALTOSID Premix** and minerals is consumed at the rate of 0.2 oz. (0.0125 lb) or 0.4 oz. (0.025 lb) per 100 lbs. of body weight per day, which is equivalent to 2 oz. or 4 oz. for a 1,000 lb. animal per day. This rate will supply the recommended average daily intake of 1.13 mg of S-methoprene per 100 lbs. of body weight per day.

MIXED-RATION FEEDING

Thoroughly mix **ALTOSID Premix** with grain or supplement. Use the following table as a guide for determining the proper mixing rate. This table provides the recommended daily intake of 1.13 mg of S-methoprene per 100 lbs. of body weight per day.

BODYWEIGHT (lbs.)	FEED RATE (lbs./day/head)	ALTOSID Premix (oz./ton)
150	5.0	6.0
200	5.0	8.0
250	5.0	10.0
300	5.0	12.0
475	5.0	19.0
700	5.0	28.0
925	5.0	37.0
1100	5.0	44.0
300	7.5	8.0
1300	15.0	17.5
1300	20.0	13.0
1300	25.0	10.5
1300	30.0	8.5
1300	35.0	7.3
1300	40.0	6.5
1300	45.0	6.0
1300	50.0	5.2
1400	20.0	14.0
1400	25.0	11.3
1400	30.0	9.1
1500	25.0	12.2
1500	30.0	9.8
1500	35.0	8.5

STORAGE AND DISPOSAL

Do not contaminate water or feed by storage or disposal.

STORAGE

Store in dry area. Do not contaminate with pesticides or fertilizer.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For information, call **1-800-347-8272** or visit our Web site: www.altosidigr.com



Wellmark International
Schaumburg, Illinois U.S.A.